

Diabetology International

Volume 4 · Number 4 · December 2013

REVIEW ARTICLE

Pathophysiology and early diagnosis of diabetic macroangiopathy

N. Katakami 205

REPORT OF THE COMMITTEE

Diagnostic criteria for acute-onset type 1 diabetes mellitus (2012) · Report of the Committee of Japan Diabetes Society on the Research of Fulminant and Acute-onset Type 1 Diabetes Mellitus

E. Kawasaki · T. Maruyama · A. Imagawa · T. Awata · H. Ikegami · Y. Uchigata · H. Osawa · Y. Kawabata · T. Kobayashi · A. Shimada · I. Shimizu · K. Takahashi · M. Nagata · H. Makino · T. Hanafusa 221

ORIGINAL ARTICLES

Combined treatment with low-dose pioglitazone and beraprost sodium improves glucose intolerance without causing body weight gain

H. Kumagai · N. Kubota · T. Kubota · T. Takahashi · M. Inoue · T. Kawai · K. Iwayama · M. Moroi · K. Sugi · T. Kadowaki 226

Evaluation of stepwise insulin treatment as an add-on to current oral antidiabetic drugs therapy in Japanese patients with type 2 diabetes: STEADY study

T. Nakagami · M. Honda · N. Shibata · O. Tomonaga · M. Tanaka · H. Kanno · J. Miura · M. Osawa · H. Takaike · Y. Iwamoto · Y. Uchigata 233

The area of abdominal subcutaneous adipose tissue is independently correlated with C-peptide increment during glucagon load in Japanese patients with type 2 diabetes

K. Matsumoto · A. Kanazawa · F. Ikeda · C. Ohmura · T. Shimizu · Y. Tamura · Y. Fujitani · R. Kawamori · H. Watada 243

Determinants of the HbA1c-lowering effect of sitagliptin when added to ongoing insulin therapy in Japanese patients with type 2 diabetes

Y. Bando · K. Ishikura · H. Kanehara · K. Aoki · A. Hisada · D. Toya · N. Tanaka 251

CASE REPORT

A diabetic patient with hemoglobin variant Hb Mito diagnosed from a change in HbA1c measurement method

M. Higa · A. Yoshifuji · A. Doi · E. Yoshida · K. Yamashita · T. Ichijo · H. Ouchi · A. Miyazaki · T. Hirose 257

SHORT COMMUNICATION

A report on diabetes seminars for medical staff: findings from a questionnaire survey

M. Kishimoto · M. Noda 261

ERRATA

Erratum to: Evaluation of a new device for measurement of hemoglobin A1c for Japanese subjects

S. Shimoda · T. Maeda · N. Furukawa · K. Ichinose · K. Taketa · M. Igata · T. Senokuchi · T. Matsumura · E. Araki 266

Erratum to: Glycated albumin as a useful clinical biomarker for glycemic variability in type 1 diabetes assessed by continuous glucose monitoring

C. Tsutsumi · A. Imagawa · M. Onishi · H. Sano · S. Nakagawa · Y. Murase-Mishiba · J. Terasaki · T. Hanafusa 267

Cover Illustration Pim-1-positive cells (*brown color*) are observed predominantly in the thickened intima of the thoracic aortas obtained at autopsy from an individual aged 35-years from the article by N. Katakami in this issue.

Further articles can be found at link.springer.com

Instructions for Authors for *Diabetol Int* are available at www.springer.com/13340